

JP42001014

Document Identifier task started on Fri Jul 21, 2006 at 4:15 PM

Explored by Document Identifier in CAPLUS and MEDLINE.

CAPLUS Answers

1 for JP42001014

MEDLINE Answers

0 for JP42001014

1 references were found (0 duplicates removed)

Connected to ChemPort: document

* Ono, Masaji, Naruse, Norio. Isoxazole derivatives. Jpn. Tokkyo Koho (1963),

Copyrights:

CAPLUS: Copyright © 2006 American Chemical Society. All Rights Reserved. (The UK patent material in this product/service is UK Crown copyright and is made available with permission. © Crown Copyright. The French (FR) patent material in this product/service is made available from Institut National de la Propriete Industrielle (INPI).)

MEDLINE: Produced by the U.S. National Library of Medicine

REGISTRY: Copyright © 2006 American Chemical Society. All Rights Reserved. (Some records contain information from GenBank(R). See also: Benson D.A., Karsch-Mizrachi I., Lipman D.J., Ostell J., Rapp B.A., Wheeler D.L. Genbank. Nucl. Acids Res. 28(1):15-18 (2000). Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.)

CASREACT: Copyright © 2006 American Chemical Society. All Rights Reserved. (In addition to reactions indexed by CAS, CASREACT contains reactions derived from the following sources: ZIC/VINITI database (1974-1991) provided by InfoChem, INPI data prior to 1986, and Biotransformations database compiled under the direction of Professor Dr. Klaus Kieslich.)

CHEMLIST, CHEMCATS: Copyright © 2006 American Chemical Society. All Rights Reserved.

JP42001014

Bibliographic Information

Isloxazole derivatives. Ono, Masaji; Naruse, Norio. (Toyo Rayon Co., Ltd.). Jpn. Tokkyo Koho (1963), 2 pp.
CODEN: JAXXAD JP 42001014 19630119 Showa. Patent written in Japanese. Application: JP 19631020. CAN
66:65459 AN 1967:65459 CAPLUS (Copyright (C) 2006 ACS on SciFinder (R))

Patent Family Information

<u>Patent No.</u>	<u>Kind</u>	<u>Date</u>	<u>Application No.</u>	<u>Date</u>
JP 42001014	B4	19630119	JP	19631020

Abstract

Manuf. of 5-aminoisoxazole derivs. by the reaction of alicyclic α -halooxime with alkali cyanide was described. In an example, 4 g. NaCN is added to 40 cc. hot (80-95°) Me₂CO, then a soln. of 10 g. α -chlorocyclooctanone oxime in 30 cc. Me₂SO dropped in during 20 min., the whole heated at 80-95° 1.5 hrs., H₂O added, and the mixt. extd. with Et₂O to give 95% 5-amino-3,4-cyclooctanoisoxazole, m. 77-8° (Et₂O-petr. ether). Similarly prepd. are 5-amino-3,4-cyclooctenoisoxazole, 5 - amino - 3,4 - trans,cis cyclododecadienoisoxazole, 5-amino-3,4 - trans,trans - cyclododecadienoisoxazole, m. 131-4°, 5-amino-3,4-cyclohexanoisoxazole (oil), and 5-amino-3,4-cyclopentaisoxazole.